

# Molecular Hydrogen Improves Obesity and Diabetes by Inducing Hepatic FGF21 and Stimulating Energy Metabolism in *db/db* Mice

Naomi Kamimura<sup>1</sup>, Kiyomi Nishimaki<sup>1</sup>, Ikuroh Ohsawa<sup>1,2</sup> and Shigeo Ohta<sup>1</sup>

Recent extensive studies have revealed that molecular hydrogen (H<sub>2</sub>) has great potential for improving oxidative stress-related diseases by inhaling H<sub>2</sub> gas, injecting saline with dissolved H<sub>2</sub>, or drinking water with dissolved H<sub>2</sub> (H<sub>2</sub>-water); however, little is known about the dynamic movement of H<sub>2</sub> in a body. First, we show that hepatic glycogen accumulates H<sub>2</sub> after oral administration of H<sub>2</sub>-water, explaining why consumption of even a small amount of H<sub>2</sub> over a short span time efficiently improves various disease models. This finding was supported by an *in vitro* experiment in which glycogen solution maintained H<sub>2</sub>. Next, we examined the benefit of *ad libitum* drinking H<sub>2</sub>-water to type 2 diabetes using *db/db* obesity model mice lacking the functional leptin receptor. Drinking H<sub>2</sub>-water reduced hepatic oxidative stress, and significantly alleviated fatty liver in *db/db* mice as well as high fat-diet-induced fatty liver in wild-type mice. Long-term drinking H<sub>2</sub>-water significantly controlled fat and body weights, despite no increase in consumption of diet and water. Moreover, drinking H<sub>2</sub>-water decreased levels of plasma glucose, insulin, and triglyceride, the effect of which on hyperglycemia was similar to diet restriction. To examine how drinking H<sub>2</sub>-water improves obesity and metabolic parameters at the molecular level, we examined gene-expression profiles, and found enhanced expression of a hepatic hormone, fibroblast growth factor 21 (FGF21), which functions to enhance fatty acid and glucose expenditure. Indeed, H<sub>2</sub> stimulated energy metabolism as measured by oxygen consumption. The present results suggest the potential benefit of H<sub>2</sub> in improving obesity, diabetes, and metabolic syndrome.

*Obesity* (2011) **19**, 1396–1403. doi:10.1038/oby.2011.6

## INTRODUCTION

Oxidative stress is involved in many lifestyle-related diseases, including diabetes, atherosclerosis, heart failure, Alzheimer's disease, and Parkinson diseases (1–6). Recent studies have revealed that molecular hydrogen (H<sub>2</sub>) acts as a novel antioxidant and prevents or ameliorates diseases associated with oxidative stress in animal experiments (7–18) and clinical tests (19–22). The brain, heart, liver, and intestine were protected from oxidative stress by inhalation of 1–2% H<sub>2</sub> gas (7–11). Interestingly, instead of inhaling H<sub>2</sub> gas, drinking water with dissolved H<sub>2</sub> (H<sub>2</sub>-water) protected the brain and kidney from oxidative stress (12–15). These studies strongly suggest the potential of H<sub>2</sub> as an effective therapeutic and preventive antioxidant; however, water dissolves H<sub>2</sub> at 0.8 mmol/l at saturated level. Thus, it has been an open question why consumption of even a small amount of H<sub>2</sub> is effective for various disease models.

Oxidative stress is one of the causes of type 2 diabetes (1–3). To examine whether H<sub>2</sub> has benefits on type 2 diabetes, we used *db/db* mice, in which oxidative stress is accumulated in

the liver and leads to hyperglycemia and hyperlipidemia (23). *Db/db* mice lack a functional leptin receptor, and have been extensively studied as a model for type 2 diabetes (24,25).

In this study, we showed that H<sub>2</sub> can be accumulated in the liver with glycogen after oral administration. Next, chronic consumption of H<sub>2</sub>-water reduced oxidative stress in the liver of *db/db* mice, and improved obesity and diabetes. As a mechanistic study, we showed that long-term consumption of H<sub>2</sub>-water enhanced the expression of a hepatic hormone, fibroblast growth factor 21 (FGF21), which is a regulator of energy expenditure (26–29). These findings suggest the great potential for hydrogen therapy and prevention of metabolic syndrome.

## METHODS AND PROCEDURES

### Animals

Male Sprague–Dawley rats of 10 weeks of age and male C57BL/6 mice of 12 weeks of age were purchased from Nippon SLC (Hamamatsu, Shizuoka, Japan). Genetically diabetic male *db/db* mice (BKS.Cg *+Lep<sup>db</sup>/+Lep<sup>db</sup>/Jcl*) and their nondiabetic heterozygous *db/+* littermates (BKS.Cg *m<sup>+</sup>/+Lep<sup>db</sup>/Jcl*) were purchased at 5 weeks of age from CLEA

<sup>1</sup>Department of Biochemistry and Cell Biology, Institute of Development and Aging Sciences, Graduate School of Medicine, Nippon Medical School, Kawasaki, Kanagawa, Japan; <sup>2</sup>Department of Environmental Gerontology, Tokyo Metropolitan Institute of Gerontology, Itabashi-ku, Tokyo, Japan. Correspondence: Shigeo Ohta (ohta@nms.ac.jp)

Received 27 July 2010; accepted 3 January 2011; published online 3 February 2011. doi:10.1038/oby.2011.6

Japan, (Tokyo, Japan). Mice were given H<sub>2</sub>-water from 6 weeks of age. For the diet-induced obesity study, C57BL/6 mice were given a high fat-diet (F2HFD1; Oriental Yeast, Tokyo, Japan) for 1 or 2 weeks. The care and use of laboratory animals were in accordance with the National Institutes of Health guidelines. This study was approved by the Animal Care and Use Committee of Nippon Medical School (Tokyo, Japan).

### Hydrogen water

H<sub>2</sub>-water was prepared as described previously (12). In brief, H<sub>2</sub> was dissolved in water under high pressure (0.4 MPa) to a supersaturated level and the saturated H<sub>2</sub>-water was stored under atmospheric pressure in an aluminum bag with no dead volume. Saturated H<sub>2</sub>-water was used as 100% H<sub>2</sub>-water. H<sub>2</sub>-water degassed by gentle stirring was used as control water. Saturated H<sub>2</sub>-water was diluted with ninefold control water and used as 10% saturated H<sub>2</sub>-water. Mice were given water freely using closed glass vessels equipped with an outlet line containing two ball bearings, which kept the water from being degassed. The vessel was freshly refilled with H<sub>2</sub>-water at 2:00 pM every day.

### Diet restriction

Diabetic mice were subjected to controlled dietary restriction, such that the daily diet consumption was equivalent to 80% or 50% of that of *ad libitum* diet consumption. Control groups were permitted free access to food and water throughout the experiment. In the diet-restricted group, mice were permitted *ad libitum* water, but food intake was restricted.

### Measurement of H<sub>2</sub> concentration

H<sub>2</sub> concentration in rat liver tissue was measured using a needle-type hydrogen electrode (Unisense, Aarhus, Denmark). Rat received H<sub>2</sub>-water orally by stomach gavage at 15 ml/kg. Throughout the experiment, the electrode current was measured with a picoammeter (Keithley, Cleveland, Ohio) and H<sub>2</sub> concentration was obtained from the calibration curve generated using known levels of H<sub>2</sub>-saturated saline.

In an *in vitro* experiment, H<sub>2</sub> concentration in glycogen, glucose solutions, or drinking water in the glass vessel was measured using a needle-type hydrogen electrode as described above.

### Sample collection and biochemical analysis

Mice were killed under anesthesia, blood was collected from the heart, and liver tissues were excised and frozen with liquid nitrogen or fixed with 4% paraformaldehyde for further analysis. The antioxidation effect of H<sub>2</sub> was determined by measuring lipid peroxides in the liver, using a malondialdehyde assay kit (Northwest Life Science Specialties, Vancouver, WA) and the level of lipid peroxides was expressed as nmol malondialdehyde (MDA)/mg protein. Plasma concentrations of total ketone bodies, triglyceride, and total cholesterol were determined with commercially available kits (Wako Pure Chemical Industries, Osaka, Japan). Plasma low-density lipoprotein-cholesterol and high-density lipoprotein-cholesterol were measured using kits (Sekisui Medical, Tokyo, Japan). Plasma glucose and nonesterified fatty acids were determined with kits available from Shino-Test (Tokyo, Japan) and Eiken Chemical (Tokyo, Japan), respectively. Plasma insulin was measured using an insulin ELISA kit (Morinaga Institute of Biological Science, Kanagawa, Japan).

### Oil Red O staining

Mouse livers were fixed in 4% paraformaldehyde in phosphate-buffered saline, embedded, and cryosectioned 10mm thick. The sections were rinsed with 60% isopropanol, stained with 0.25% Oil Red O solution, rinsed with 60% isopropanol, and mounted in aqueous mountant. The area of stained lipid (%) was calculated using the Image J program (ver 1.41; National Institutes of Health, Bethesda, MD) from four sections for each mouse.

### Body fat composition analysis

For computed tomography analysis of body fat composition, mice were anesthetized with halothane in a mixture of nitrous oxide and oxygen (70%:30%, vol/vol) and then scanned using a LaTheta LCT-100,

experimental animal computed tomography system (Aloka, Tokyo, Japan). Contiguous 1-mm slice images were used for quantitative assessment using LaTheta software (ver 1.00). Visceral fat, subcutaneous fat, and muscle were distinguished and evaluated quantitatively.

### RNA isolation and reverse transcriptase-PCR

Total RNA was isolated from the liver tissue using an RNeasy Mini kit (QIAGEN, Valencia, CA). Complementary DNA generated by SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, CA) was analyzed by quantitative PCR using Thermal Cycler Dice Real Time System TP800 (TAKARA BIO, Shiga, Japan). All samples were normalized to glyceraldehyde 3-phosphate dehydrogenase expression. Primer and probe sequences for each PCR are shown in Table 1.

### Behavioral analysis

Movement activities were recorded in *db/+* and *db/db* mice automatically using a laboratory animal monitoring system (ACTIMO-100; Shinfactory, Fukuoka, Japan). Mice were housed individually and temperature was maintained at 22 °C. Food and water with or without hydrogen were available *ad libitum*. Mice were acclimatized to the chambers for 24h before beginning recordings and then monitored for 48h. Movement activity was measured as ambulatory counts from a record of consecutive adjacent infrared beam breaks. Cumulative ambulatory counts on the x- and y-axes were recorded every 10 min.

### Indirect calorimetric analyses

Metabolic rate was measured by indirect calorimetric analysis in *db/+* and *db/db* mice using an open-circuit calorimeter (Oxymax; Columbus Instruments, Columbus, OH). Mice were housed individually in a chamber (20 × 10 × 12.7 cm) and temperature was maintained at 22 °C, with air flow of 0.5l/min. Food and water with or without H<sub>2</sub> were available *ad libitum*. Mice were acclimatized to the chambers for 48h before beginning recordings and then monitored for 48h. VO<sub>2</sub> and VCO<sub>2</sub> were measured every 10 min using an electrochemical O<sub>2</sub> analyzer and a CO<sub>2</sub> sensor (Oxymax), and the respiratory exchange ratio was calculated as VCO<sub>2</sub>/VO<sub>2</sub> (volume of CO<sub>2</sub> produced per volume of O<sub>2</sub> consumed (ml/kg/h)).

### Statistical analysis

We performed statistical analysis using StatView software (SAS Institute) by applying an unpaired two-tailed Student's *t*-test and ANOVA followed by Fisher's exact test, as described previously (7). Differences were considered statistically significant at *P* < 0.05.

**Table 1 Primers and probes for reverse transcriptase-PCR**

Gene		Sequence
FGF21	F primer	5'-CCGCAGTCAGAAAGTCTCCT-3'
	R primer	5'-TCTGAAGCTGCAGGCCTCA-3'
	Probe	5'-AGCTCTCTATGGATCGCCTCACTTTGATCC-3'
PEPCK	F primer	5'-TGCTGCAGAACACAAGGGC-3'
	R primer	5'-TTTGCCGAAGTTGTAGCCG-3'
	Probe	5'-TCATCATGCACGACCCCTTTGCC-3'
G6PC	F primer	5'-CGCCATGCAAAGGACTAGGA-3'
	R primer	5'-AGGGCCGATGTCAACACCT-3'
	Probe	5'-TAAAGCCTCTGAAACCCATTGTGAGGCC-3'
GAPDH	F primer	5'-CATCACTGCCACCCAGAAGA-3'
	R primer	5'-ATGTTCTGGGCAGCC-3'
	Probe	5'-TGGATGGCCCCCTCTGGAAGCTG-3'


FGF21, fibroblast growth factor 21; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; G6PC, glucose-6-phosphatase, catalytic subunit; PEPCK, phosphoenolpyruvate carboxykinase.

## RESULTS

**Molecular hydrogen is accumulated in the liver with glycogen**

We monitored the dynamic movement of  $H_2$  in the liver after oral administration of  $H_2$ -water. Rat received  $H_2$ -water orally by stomach gavage, and levels of  $H_2$  in the liver were monitored by directly inserting a needle-type hydrogen sensor into the liver for an hour (Figure 1a). The  $H_2$  concentration profile gave a peak 5 min after administration of  $H_2$ -water in both a fed and fasted liver; however, a great difference was found between the fed and fasted liver; the maximum  $H_2$  level in the fed liver was twofold that of the fasted liver. Moreover, the fed liver maintained a considerable  $H_2$  level for an hour, while that in the fasted liver returned to the basal level after 25 min.

Since a fed liver is rich in glycogen, we speculated that higher polymers of carbohydrates including glycogen have the capacity to maintain  $H_2$ . To verify this speculation, we examined saturated solubilities of  $H_2$  in glycogen and glucose solutions by a hydrogen sensor.  $H_2$  was dissolved in a glycogen or glucose solution by bubbling  $H_2$  gas up to a saturated level (Figure 1b). Compared to the glucose solution and water, the glycogen solution dissolved a significantly higher amount of



**Figure 1** Hydrogen is accumulated and maintained in a fed liver and in glycogen solution *in vitro*. (a) The concentration of molecular hydrogen in the liver was monitored using a needle-type hydrogen sensor in fed or overnight-fasted rat liver. Rat received hydrogen water (0.8 mmol/l  $H_2$  in water) orally by stomach gavage at 15 ml/kg. Arrow indicates the time point when rat was administered hydrogen water. (b) Saturated concentration of hydrogen in glycogen and glucose solutions, and water. Molecular hydrogen was dissolved in indicated solutions by bubbling  $H_2$  gas to a saturated level. The concentration of hydrogen in solution was measured using a hydrogen sensor. Data are means  $\pm$  s.d. ( $n = 3$ ). \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$  compared with 100 or 10 mg/ml glycogen groups. (c) Concentration of hydrogen in water, 100 mg/ml glycogen solution and 100 mg/ml glucose solution after bubbling stopped. Hydrogen-saturated solutions were kept in a plastic tube with the lid off at 20 °C for 0, 30, or 90 min under atmospheric pressure. Concentration of hydrogen in solution was measured using a hydrogen sensor. Data are mean  $\pm$  s.d. ( $n = 3$ ). \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .

$H_2$ . The hydrogen sensor was not influenced by glycogen and glucose (data not shown). In addition,  $H_2$  was maintained for a longer time in saturated glycogen solution than in saturated glucose solution or water in a plastic tube with the lid off under atmospheric pressure (Figure 1c). The half-life was  $19.1 \pm 2.3$ ,  $30.9 \pm 3.0$ , and  $20.3 \pm 0.4$  min (mean  $\pm$  s.d.,  $n = 3$ ) in water, 100 mg/ml glycogen and glucose solutions, respectively. The half-life of dissolving  $H_2$  in the glycogen solution was prolonged 1.6-fold. Thus, it is concluded that  $H_2$  can be accumulated and reserved in the liver with glycogen, suggesting that expenditure of glycogen should accompany the release of  $H_2$ .


**Consuming hydrogen water reduces oxidative stress in the liver and improves fatty liver**

Since obesity is a proinflammatory disease, consumption of  $H_2$ -water may suppress obesity by acting as an anti-inflammatory. To examine the antioxidation effect of  $H_2$  on the liver, we used obesity and type 2 diabetes model mice *db/db* lacking functional leptin receptors, because oxidative stress is accumulated in the liver. *Db/db* mice and their lean littermates drank  $H_2$ -water *ad libitum* for 3 months.  $H_2$ -water was exchanged for fresh saturated or 10% saturated water at 1400 h every day. The concentration of dissolved  $H_2$  was measured as described in Methods and Procedures (Figure 2a). Since *db/db* mice drank much more water than wild-type mice,  $H_2$  was degassed much faster into the air phase in the vessel used for *db/db* mice; however, a considerable amount of  $H_2$  was maintained by this method.

The effect of  $H_2$  on oxidative stress in the liver was examined as judged by the level of malondialdehyde (MDA), an oxidative stress marker derived from lipid peroxides. The MDA level in the liver of  $H_2$ -administered mice significantly fell to nearly the level in nondiabetes control mice, indicating that consumption of  $H_2$ -water *ad libitum* markedly suppressed oxidative stress (Figure 2b). A dotted pattern caused by the accumulation of fat disappeared in the liver of *db/db* mice with  $H_2$ , compared to control water administered *db/db* mice (Figure 2c, insets of upper panels). Oil red O staining indicated that  $H_2$  administration significantly reduced neutral lipid accumulation in the livers of *db/db* mice (Figure 2b, lower panels and Figure 2d, left graph). Furthermore, even with short-term administration (1–2 weeks),  $H_2$  significantly reduced fat accumulation in the liver of high fat-diet-induced obesity mice using the wild-type (Figure 2d, middle and right graph). These data clearly indicate that consumption of  $H_2$  markedly reduces hepatic oxidative stress levels and improves fatty liver in *db/db* as well as diet-induced obesity mice.

**Consuming hydrogen water suppressed body-weight gain and reduced plasma glucose and triglyceride levels**

To investigate the effect of  $H_2$  on the obesity of *db/db* mice, body-weight was monitored throughout the experimental period, and body fat mass at 18 weeks old was measured by computerized tomography. Mice were divided randomly into 3 groups. Group I (control) was allowed to freely drink water



**Figure 2** Consuming hydrogen suppresses oxidative stress in the liver and improves fatty liver. **(a)** Hydrogen concentration in a glass vessel described in Methods and Procedures section was measured without mice (open circle), showing that the equipment retains hydrogen. Profile of the hydrogen concentration in drinking water in a glass vessel given to *db/db* mice (closed square) or wild-type mice used for diet-induced obesity (DIO) (closed circle), indicating that hydrogen gas was released into the air phase by consuming water. Data are the mean  $\pm$  s.d. ( $n = 3$ ). **(b)** Malondialdehyde concentration in a fed or overnight-fasted liver was measured. The *db/+* and *db/db* mice were given water with or without hydrogen for 3 months. Data are mean  $\pm$  s.e.m. ( $n = 10$  for each *db/+* group and  $n = 15$  for each *db/db* group). \* $P < 0.05$ , \*\* $P < 0.01$ . **(c)** The appearance and representative oil red O staining of the liver of *db/+* and *db/db* mice given water with and without hydrogen for 3 months, respectively. Scale bar: 100  $\mu$ m. **(d)** Levels of fat accumulation in the liver. The *db/db* mice were given water with or without hydrogen for 3 months (left panel). DIO (high fat-diet induced obesity) mice were given water with or without hydrogen for 1 or 2 weeks (middle or right panel). Oil accumulation in the liver was calculated from oil red O staining using an image analysis program, Image J program. Data are mean  $\pm$  s.e.m. ( $n = 12$  for each *db/db* group and  $n = 8$  for each DIO group). \* $P < 0.05$ , \*\* $P < 0.01$ .

without H<sub>2</sub>. Group II and Group III were given H<sub>2</sub>-water with 0.8 mmol/l (saturated H<sub>2</sub>-water; 100%) and 0.08 mmol/l (10% saturated level of H<sub>2</sub>), respectively. In the initial phase of the experiment, the weight gains of all group animals were similar, suggesting no toxic effect of H<sub>2</sub>-water; however, while the control mice exhibited a progressive weight increase, the mice in both 100% and 10% H<sub>2</sub>-administered groups achieved a modest weight increase (Figure 3b). Group II mice (100% H<sub>2</sub>) were slimmer and their body-weight was significantly lower than that of Group I mice (control) at 18 weeks of age (Figure 3a,b). The suppression of body-weight gain was observed in mice drinking 10% H<sub>2</sub>-water (Group III) (Figure 3b). Body fat was also substantially lower in 100% H<sub>2</sub>-consuming mice (Figure 3c,d). Since the consumed amounts and volumes of diet and water did not differ among groups (Figure 3e,f), it is suggested that H<sub>2</sub> consumption stimulates energy metabolism to suppress the gain of fat and body weights.


Next, we performed biochemical examinations of blood. Plasma levels of glucose and insulin were significantly reduced in the 100% H<sub>2</sub>-administered group and triglycerides were significantly lowered in both 100% and 10% H<sub>2</sub>-administered groups (Figure 4a–c). Plasma total ketone bodies tended to increase in the 100% H<sub>2</sub>-administered group (Figure 4d), while no change in plasma levels of free fatty acid, high-density lipoprotein-cholesterol, low-density lipoprotein-cholesterol, total cholesterol, and adiponectin was found (Figure 4e–i). These data demonstrate that the consumption of H<sub>2</sub> markedly improves obesity, hyperglycemia, and the plasma triglycerides of diabetic *db/db* mice.

### Consuming hydrogen water shows a similar effect to diet restriction

Since *db/db* mice cannot regulate their appetite to eat excess diet by a deficient leptin receptor gene, diet restriction should be effective to improve obesity and diabetes (30). When *db/db* mice were subjected to controlled dietary restriction, the levels of plasma glucose, insulin, and triglycerides significantly fell (Figure 5a–c). When comparing the effects by drinking H<sub>2</sub>-water and dietary restriction, the plasma glucose level of mice given 80% of *ad libitum* diet consumption was the same as that of H<sub>2</sub>-administered mice (Figure 5a). Furthermore, an additive effect was observed when mice were given both H<sub>2</sub>-water and a restricted diet (Figure 5a). A similar additive effect was seen in plasma triglyceride levels (Figure 5c). In particular, when both H<sub>2</sub>-water and 80% diet restriction were given to *db/db* mice, the plasma triglyceride reached the level of control *db/+* mice (Figure 5c).

### Consuming hydrogen water increases hepatic mRNA level of FGF21

The liver plays an essential role in controlling blood glucose levels by modulating glucose catabolism and gluconeogenesis. To understand the mechanism of how H<sub>2</sub> regulates glucose and triglycerides, *db/db* mice given H<sub>2</sub>-water for 3 months were subjected to expression analyses of genes related with the regulation of gluconeogenesis. Phosphoenolpyruvate carboxykinase catalyzes the initial step of gluconeogenesis, and glucose-6-phosphatase catalyzes the last committed step of this process (31). Data from real-time PCR revealed that H<sub>2</sub> administration had no influence on hepatic mRNA levels of



**Figure 3** Consuming molecular hydrogen suppresses obesity. (a) The appearance of *db/db* mice given water with or without hydrogen for 3 months. Each average mouse was photographed. (b) Body weights of *db/db* mice given water with 100% (0.8 mmol/l) or 10% (0.08 mmol/l) hydrogen were examined every 2 weeks. Control mice (CTL) received water without hydrogen. \* $P < 0.05$ ; 100%  $H_2$  vs. control group ( $n = 9$  for  $H_2$  and  $n = 6$  for control group). (c) Representative CT images of abdominal part of average *db/db* mice drinking water with or without hydrogen for 3 months. Blue, green, and red represent visceral fat, subcutaneous fat, and muscle, respectively. (d) Total body fat composition of *db/db* mice calculated by the integration of fat area in each section from CT scan images. Data are mean  $\pm$  s.e.m. ( $n = 15$ ). \* $P < 0.05$ . (e) Food and (f) water intake of *db/db* mice was measured every 2 weeks throughout the experiment.


phosphoenolpyruvate carboxykinase and glucose-6-phosphatase, catalytic subunit (G6PC) (Figure 6c,d).

Since FGF21, an atypical member of the fibroblast growth factor family, contributes to energy metabolism, we focused on the gene expression of FGF21 after integrated study of the gene expression.  $H_2$  administration induced hepatic mRNA levels of FGF21, regardless of a fed or fasted liver (Figure 6a,b). These results indicate that, at least in part, the induction of hepatic FGF21 contributes to the lowering effect on plasma glucose and triglyceride levels.

#### Drinking hydrogen water stimulates energy metabolism

To verify whether drinking  $H_2$ -water stimulates energy metabolism, we examined oxygen ( $O_2$ ) consumption and carbon dioxide ( $CO_2$ ) production. First, we compared physical ability by detecting the movement of mice with infrared beams. Although *db/db* mice was apparently less active in light and dark than control *db/+* mice, no difference was found between *db/db* mice with and without  $H_2$ -water (Figure 7a,b).

*Db/db* mice with or without  $H_2$ -water consumed less  $O_2$  and produced less  $CO_2$  than *db/+* control mice (Figure 7c-f).




**Figure 4** Consuming molecular hydrogen suppresses hyperglycemia, hyperinsulinemia, and plasma triglyceride level. *Db/db* mice were given water with 100% (0.8 mmol/l) or 10% (0.08 mmol/l) hydrogen for 3 months. *Db/+* and *db/db* mice were given water without hydrogen (HW) for the same period as controls. Biochemical analyses were performed to obtain plasma parameters of *db/+* and *db/db* mice. (a) Plasma concentration of glucose, (b) insulin, (c) triglyceride (TG), (d) total ketone bodies (T-KB), (e) free fatty acids (NEFA), (f) high-density lipoprotein-cholesterol (HDL-C), (g) low-density lipoprotein-cholesterol (LDL-C), (h) total cholesterol (T-CHO), and (i) adiponectin (ADN) are shown as mean  $\pm$  s.e.m. ( $n = 15$ ). \* $P < 0.05$ , \*\* $P < 0.01$ .

Moreover, it is significant that  $H_2$ -drinking *db/db* mice consumed more  $O_2$ , 10%, and produced more  $CO_2$ , 10%, than *db/db* mice without  $H_2$ -water during both night and day (Figure 7c-f). Since respiratory exchange rates ( $VCO_2/VO_2$ ) did not differ between *db/db* mice with and without  $H_2$ -water, the carbon source for energy production was not changed to stimulate energy metabolism (Figure 7g,h).


Thus, drinking  $H_2$ -water suppresses the gain of fat and body weights and improves metabolic parameters by stimulating energy metabolism.

#### DISCUSSION


In this study, we show a novel benefit of  $H_2$ , it may be useful on therapy for and prevention of obesity and diabetes. So far, many reports have confirmed that consumption of  $H_2$  reduces oxidative stress in various disease models and clinical tests (7,8,10-16,19). Clinical tests revealed that drinking  $H_2$ -water reduced oxidative stress makers in patients with type 2 diabetes (19) or subjects with potential metabolic syndrome (20), and influenced glucose (19) and cholesterol metabolism (20).



**Figure 5** The effect of hydrogen is similar to that of dietary restriction. *Db/db* mice were given water with or without hydrogen (HW) and subjected to dietary restriction (DR). Control groups were permitted free access to food and water throughout the experiment. In diet-restricted groups, mice were permitted *ad libitum* water, but food intake was restricted to 80% or 50% of that of *ad libitum* diet consumption. Plasma concentrations of (a) glucose, (b) insulin (INS), and (c) triglyceride (TG) were measured after 3-month treatment. Data are mean  $\pm$  s.e.m. \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ ; compared with HW (-)/DR (-) group ( $n = 8-15$ ).



**Figure 7** Molecular hydrogen increases oxygen consumption and carbon dioxide production without influencing movement activities. (a and b) movement activity, (c and d) oxygen consumption, (e and f) carbon dioxide production, and (g and h) RER (respiratory exchanging rate) in *db/+* and *db/db* mice given water with or without 100% (0.8 mmol/l) hydrogen for 3 months. (a, c, e, and g) Representative profiles for each parameter. Gray area represent dark phase. (b, d, f, and h) Data are the mean  $\pm$  s.e.m. ( $n = 9$  and  $n = 6$ , for *db/db* and for *db/+* groups, respectively). \* $P < 0.05$ .



**Figure 6** Chronic effects of hydrogen on fibroblast growth factor 21 (FGF21) gene expression in the liver. *Db/+* and *db/db* mice were given water with or without hydrogen for 3 months. (a) The gene expression of FGF21 with fed liver, (b) FGF21 with fast liver, (c) Phosphoenolpyruvate carboxykinase (PEPCK), and (d) glucose-6-phosphatase, catalytic subunit (G6PC) were measured. Data are mean  $\pm$  s.e.m. \* $P < 0.05$ ; control vs. hydrogen water in *db/db* mice ( $n = 15$ ).

The mitochondria are a major source of reactive oxygen species during energy production metabolism and  $H_2$  directly protects mitochondria that are exposed to reactive oxygen species (7). Thus, it may be reasonable that mitochondrial energy metabolism, especially fatty acid metabolism, functions against oxidative stress to efficiently expend glucose and fatty acid.

Initially,  $H_2$  was dissolved in culture media and shown to protect cells and organelles by directly reacting highly active reactive oxygen species (7). Next, the brain, heart, liver, and intestine were protected from oxidative stress by inhalation of  $H_2$  gas (7-11). Interestingly, instead of inhaling  $H_2$  gas,  $H_2$ -water was effective in protecting the brain and kidney from oxidative stress (12-15) and decreased oxidative stress markers of patients with diabetes and potential metabolic syndrome (19,20). Drinking  $H_2$ -water is the most convenient method to

consume H<sub>2</sub>; however, when H<sub>2</sub> is consumed from H<sub>2</sub>-water, a very limited amount of H<sub>2</sub> can be consumed, because H<sub>2</sub> is dissolved in water at a saturated level of only 0.8 mmol/l under atmospheric pressure. After H<sub>2</sub>-water is consumed, most H<sub>2</sub> in blood becomes undetectable within 30 min (14), probably *via* expiration from the lungs; thus, it has been an open question why H<sub>2</sub>-water is effective despite its small amount and short exposure. In this study, it was found that H<sub>2</sub> can be accumulated and reserved in the liver with glycogen, which at least partly explains this question.

Moreover, it has been reported that H<sub>2</sub> acts as an anti-inflammatory and antiallergic regulator by inducing inflammatory cytokines, such as tumor necrosis factor- $\alpha$ , interleukin-6 and some phosphorylating signal factors (9,10,14,32). On the other hand, the concept that obesity is a proinflammatory disease has been accepted (33); thus, consumption of H<sub>2</sub>-water may suppress obesity by acting as an anti-inflammatory.

Alternatively, we found that consumption of H<sub>2</sub>-water enhanced the expression of FGF21. Since it has not been shown that H<sub>2</sub> directly regulates transcription, H<sub>2</sub> is indirectly involved in FGF21 expression. Since FGF21 is a metabolic hormone that improves insulin sensitivity and glucose clearance, reduces plasma triglyceride concentrations and suppresses weight gain when fed a high-fat diet (26–29), all the findings shown in this study were elucidated by the enhanced expression of FGF21.

Indeed, we revealed that drinking H<sub>2</sub>-water stimulates energy metabolism as measured by O<sub>2</sub> consumption and CO<sub>2</sub> production. The enhancement of energy metabolism may fully elucidate why consumption of H<sub>2</sub>-water suppresses the gain of fat and body weights and improves metabolic parameters; however, it remains unknown whether the induction of FGF21 fully elucidates the enhancement of energy metabolism. Moreover, the relationships among the reduction of oxidative stress, induction of FGF21 expression and stimulation of energy metabolism are still unclear. It should be analyzed whether these relationships are direct or indirect. It might be valuable to examine the relationship of obesity with oxidative stress. Although the primary target of H<sub>2</sub> essentially remains unknown, these findings provide a clue to understand the mechanism of chronic treatment with H<sub>2</sub>-water.

Finally, we would like to emphasize that the novel benefit of H<sub>2</sub> in therapeutic and preventive applications for metabolic syndrome could be achieved by the most convenient way, H<sub>2</sub>-water.

#### ACKNOWLEDGMENTS

We thank Blue Mercury (Tokyo, Japan) for the generous gift of hydrogen water and M. Takeda for technical assistance. This work was supported by a grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No.20300230 for S.O. and No.20500723 for N.K.) and the Ministry of Health, Labour and Welfare for S.O. (Intramural Research Grant (20B-13) for Neurological and Psychiatric Disorders of NCNP).

#### DISCLOSURE

The authors declared no conflict of interest.

#### REFERENCES

- Pérez-Matute P, Zulet MA, Martínez JA. Reactive species and diabetes: counteracting oxidative stress to improve health. *Curr Opin Pharmacol* 2009;9:771–779.
- Roberts CK, Sindhu KK. Oxidative stress and metabolic syndrome. *Life Sci* 2009;84:705–712.
- Reddy VP, Zhu X, Perry G, Smith MA. Oxidative stress in diabetes and Alzheimer's disease. *J Alzheimers Dis* 2009;16:763–774.
- Fearon IM, Faux SP. Oxidative stress and cardiovascular disease: novel tools give (free) radical insight. *J Mol Cell Cardiol* 2009;47:372–381.
- Tsang AH, Chung KK. Oxidative and nitrosative stress in Parkinson's disease. *Biochim Biophys Acta* 2009;1792:643–650.
- Henchcliffe C, Beal MF. Mitochondrial biology and oxidative stress in Parkinson disease pathogenesis. *Nat Clin Pract Neurol* 2008;4:600–609.
- Ohsawa I, Ishikawa M, Takahashi K *et al*. Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals. *Nat Med* 2007;13:688–694.
- Fukuda K, Asoh S, Ishikawa M *et al*. Inhalation of hydrogen gas suppresses hepatic injury caused by ischemia/reperfusion through reducing oxidative stress. *Biochem Biophys Res Commun* 2007;361:670–674.
- Nakao A, Kaczorowski DJ, Wang Y *et al*. Amelioration of rat cardiac cold ischemia/reperfusion injury with inhaled hydrogen or carbon monoxide, or both. *J Heart Lung Transplant* 2010;29:544–553.
- Buchholz BM, Kaczorowski DJ, Sugimoto R *et al*. Hydrogen inhalation ameliorates oxidative stress in transplantation induced intestinal graft injury. *Am J Transplant* 2008;8:2015–2024.
- Hayashida K, Sano M, Ohsawa I *et al*. Inhalation of hydrogen gas reduces infarct size in the rat model of myocardial ischemia-reperfusion injury. *Biochem Biophys Res Commun* 2008;373:30–35.
- Nagata K, Nakashima-Kamimura N, Mikami T, Ohsawa I, Ohta S. Consumption of molecular hydrogen prevents the stress-induced impairments in hippocampus-dependent learning tasks during chronic physical restraint in mice. *Neuropsychopharmacology* 2009;34:501–508.
- Nakashima-Kamimura N, Mori T, Ohsawa I, Asoh S, Ohta S. Molecular hydrogen alleviates nephrotoxicity induced by an anti-cancer drug cisplatin without compromising anti-tumor activity in mice. *Cancer Chemother Pharmacol* 2009;64:753–761.
- Cardinal JS, Zhan J, Wang Y *et al*. Oral hydrogen water prevents chronic allograft nephropathy in rats. *Kidney Int* 2010;77:101–109.
- Fujita K, Seike T, Yutsudo N *et al*. Hydrogen in drinking water reduces dopaminergic neuronal loss in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease. *PLoS ONE* 2009;4:e7247.
- Fu Y, Ito M, Fujita Y *et al*. Molecular hydrogen is protective against 6-hydroxydopamine-induced nigrostriatal degeneration in a rat model of Parkinson's disease. *Neurosci Lett* 2009;453:81–85.
- Ohsawa I, Nishimaki K, Yamagata K, Ishikawa M, Ohta S. Consumption of hydrogen water prevents atherosclerosis in apolipoprotein E knockout mice. *Biochem Biophys Res Commun* 2008;377:1195–1198.
- Ohazawa H, Igarashi T, Yokota T *et al*. Protection of the retina by rapid diffusion of hydrogen: administration of hydrogen-loaded eye drops in retinal ischemia-reperfusion injury. *Invest Ophthalmol Vis Sci* 2010;51:487–492.
- Kajiyama S, Hasegawa G, Asano M *et al*. Supplementation of hydrogen-rich water improves lipid and glucose metabolism in patients with type 2 diabetes or impaired glucose tolerance. *Nutr Res* 2008;28:137–143.
- Nakao A, Toyoda Y, Sharma P, Evans M, Guthrie N. Effectiveness of hydrogen rich water on antioxidant status of subjects with potential metabolic syndrome—an open label pilot study. *J Clin Biochem Nutr* 2010;46:140–149.
- Nakayama M, Kabayama S, Nakano H *et al*. Biological effects of electrolyzed water in hemodialysis. *Nephron Clin Pract* 2009;112:c9–15.
- Suzuki Y, Sano M, Hayashida K *et al*. Are the effects of  $\alpha$ -glucosidase inhibitors on cardiovascular events related to elevated levels of hydrogen gas in the gastrointestinal tract? *FEBS Lett* 2009;583:2157–2159.
- Kumashiro N, Tamura Y, Uchida T *et al*. Impact of oxidative stress and peroxisome proliferator-activated receptor  $\gamma$  coactivator-1 $\alpha$  in hepatic insulin resistance. *Diabetes* 2008;57:2083–2091.
- Coleman DL. Obese and diabetes: two mutant genes causing diabetes-obesity syndromes in mice. *Diabetologia* 1978;14:141–148.
- Leibel RL, Chung WK, Chua SC Jr. The molecular genetics of rodent single gene obesities. *J Biol Chem* 1997;272:31937–31940.

26. Reitman ML. FGF21: a missing link in the biology of fasting. *Cell Metab* 2007;5:405–407.
27. Kharitonov A, Shiyanova TL, Koester A *et al*. FGF-21 as a novel metabolic regulator. *J Clin Invest* 2005;115:1627–1635.
28. Kharitonov A, Wroblewski VJ, Koester A *et al*. The metabolic state of diabetic monkeys is regulated by fibroblast growth factor-21. *Endocrinology* 2007;148:774–781.
29. Kharitonov A, Shanafelt AB. FGF21: a novel prospect for the treatment of metabolic diseases. *Curr Opin Investig Drugs* 2009;10:359–364.
30. Lee SM, Bressler R. Prevention of diabetic nephropathy by diet control in the db/db mouse. *Diabetes* 1981;30:106–111.
31. Ruderman NB. Muscle amino acid metabolism and gluconeogenesis. *Annu Rev Med* 1975;26:245–258.
32. Itoh T, Fujita Y, Ito M *et al*. Molecular hydrogen suppresses FcepsilonRI-mediated signal transduction and prevents degranulation of mast cells. *Biochem Biophys Res Commun* 2009;389:651–656.
33. Aggarwal BB. Targeting inflammation-induced obesity and metabolic diseases by curcumin and other nutraceuticals. *Annu Rev Nutr* 2010;30:173–199.